

# Dystan<sup>®</sup>-Powder

agilan<sup>®</sup> complementary feed  
with spruce needles, wormwood, balm, chamomile and  
pure apple pectin

For all animal species

## Composition

Dextrose  
Spruce needles (11 %)  
Apple pectin (10 %)  
Sodium chloride  
Powdered milk  
Psyllium husks ground  
Sodium bicarbonate  
Potassium chloride

## Analytical constituents

Crude protein	11.7 %
Crude fat	2.9 %
Crude fibre	10.1 %
Crude ash	20.5 %
Sodium	4.0 %
Lysine	0.2 %
Methionine	0.0 %

## Content of additives per kg

### Trace elements:

Copper (3b405) (as Cu-II-sulphate, pentahydrate)	750 mg
Zinc (3b605) (as Zn sulphate, monohydrate)	4,000 mg
Manganese (3b503) (as manganous-(II)-sulphate, monohydrate)	3,000 mg

### Flavouring compounds:

Melissa officinalis 70/524/EEC; CoE 280 (Extract)	4,000 mg
Matricaria recutita 70/524/EEC (Extract)	5,000 mg
Artemisia absinthium L. 70/524/EEC (Tincture)	1,000 mg

### Acidity regulator:

Citric acid (1a330)	78.000 mg
---------------------	-----------

## Recommended use

With natural ingredients such as lemon balm, chamomile and wormwood for supplementary feeding as needed for digestive disorders and intestinal problems such as a change in faecal consistency, flatulence and a general reluctance to eat.

## Feeding recommendation

The present statements apply as long as the veterinarian has not advised otherwise:



# Dystan®-Powder

- Calves:  
As needed, 2-3 x a day dissolve 50 g Dystan® Powder in 1 - 2 litres of warm water (40 °C) and add to feed (or 0.1 – 1.0 g/ kg BW [body weight]).
- Cattle, horses:  
50 - 75 g per administration (or 0.1 g/ kg body weight)
- Foals:  
up to 25 g per administration (or 0.1 - 0.5 g/ kg body weight)
- Pigs (weaners, shoats):  
up to 15 g per administration (or 0.1 - 0.5 g/ kg body weight)
- Sucklings  
up to 6 g per administration (or 0.1 - 1.0 g/ kg body weight)
- Dogs, sheep:  
up to 25 g per administration (or 0.1 - 1.0 g/ kg body weight)
- Poultry:  
0.1 - 1.0 g/ kg body weight  
(do not administer through drinking water system)
- Small domestic animals:  
0.1 - 1.0 g/ kg body weight

**Do not use this product as the exclusive source of drinking water!**

## Feeding advice

The simultaneous use of different organic acids or their salts is contraindicated where one or more of them is used at or near the maximum permitted content.

## Storage

Store in a dry place at room temperature.  
Close package tightly after use so it cannot absorb any moisture.  
Shelf life guarantee only applies to unopened containers.

Date of revision of the text: February 2024